

Comparison of peripapillary retinal nerve fiber layer thickness between myopia severity groups and controls

Evelyn Li Min Tai^{1,2}, Jiunn Loong Ling^{1,2}, Eng Hui Gan^{1,2}, Hussein Adil^{1,2}, Wan-Hitam Wan-Hazabbah^{1,2}

¹Department of Ophthalmology, School of Medical Sciences, Health Campus, Universiti Sains Malaysia, Kubang Kerian 16150, Kelantan, Malaysia

²Hospital Universiti Sains Malaysia, Kubang Kerian 16150, Kelantan, Malaysia

Correspondence to: Evelyn Li Min Tai. Department of Ophthalmology, School of Medical Sciences, Health Campus, Universiti Sains Malaysia, Kubang Kerian 16150, Kelantan, Malaysia. daileid@yahoo.com

Received: 2017-05-12 Accepted: 2017-10-23

Abstract

• **AIM:** To compare the peripapillary retinal nerve fiber layer (RNFL) thickness measured *via* optical coherence tomography (OCT) between different groups of myopia severity and controls.

• **METHODS:** This was a prospective cross-sectional study. All subjects underwent a full ophthalmic examination, refraction, visual field analysis and A-scan biometry. Myopic patients were classified as low myopia (LM) [spherical equivalent (SE) from greater than -0.5 D, up to -3.0 D], moderate myopia (MM; SE greater than -3.0 D, up to -6.0 D) and high myopia (HM; SE greater than -6.0 D). The control group consisted of emmetropic (EM) patients (SE from +0.5 D to -0.5 D). A Zeiss Cirrus HD-OCT machine was used to measure the peripapillary RNFL thickness of both eyes of each subject. The mean peripapillary RNFL thickness between groups was compared using both analysis of variance and analysis of covariance.

• **RESULTS:** A total of 403 eyes of 403 subjects were included in this study. The mean age was 31.48±10.23y. There were 180 (44.7%) eyes with EM, 124 (30.8%) with LM, 73 (18.1%) with MM and 26 (6.5%) with HM. All groups of myopia severity had a thinner average RNFL than the EM group, but after controlling for gender, age, and axial eye length, only the HM group differed significantly from the EM group ($P=0.017$). Likewise, the superior, inferior and nasal RNFL was thinner in all myopia groups compared to controls, but after controlling for confounders, only the inferior quadrant RNFL was significantly thinner in the HM group, when compared to the EM group ($P=0.017$).

• **CONCLUSION:** The average and inferior quadrant RNFL is thinner in highly myopic eyes compared to emmetropic eyes. Refractive status must be taken into consideration

when interpreting the OCT of myopic patients, as RNFL thickness varies with the degree of myopia.

• **KEYWORDS:** optical coherence tomography; retinal nerve fiber layer; myopia

DOI:10.18240/ijo.2018.02.16

Citation: Tai ELM, Ling JL, Gan EH, Adil H, Wan-Hazabbah WH. Comparison of peripapillary retinal nerve fiber layer thickness between myopia severity groups and controls. *Int J Ophthalmol* 2018;11(2):274-278

INTRODUCTION

The prevalence of myopia has been increasing over the past decades, with a projected half of the world population estimated to be myopic by 2050^[1-2]. This condition is especially common in East Asia, where the prevalence has been estimated to be as high as 90%^[3-4]. Myopia has been associated with an increased risk of various other eye diseases, of which glaucoma remains one of the pertinent^[5-6].

The morphological appearance of the optic nerve head in myopia renders the clinical diagnosis and monitoring of glaucoma progression in myopic eyes challenging, especially as these eyes may have concomitant visual field defects mimicking those seen in glaucoma^[7]. Imaging modalities such as optical coherence tomography (OCT) can aid in the diagnostic dilemma by measuring retinal nerve fiber layer (RNFL) thickness, which differs significantly between glaucoma patients and controls^[8-11]. However, myopic patients may have RNFL abnormalities which may complicate this interpretation. Our study thus aimed to compare peripapillary RNFL thickness between different groups of myopia severity and controls using Zeiss Cirrus HD-OCT.

SUBJECTS AND METHODS

This was a prospective cross-sectional study conducted in the Eye Clinic of Hospital Universiti Sains Malaysia (USM), a tertiary hospital in Malaysia. The study was approved by the Human Research Ethics Committee of USM, and the conduct of the study followed the tenets of the Declaration of Helsinki. Inclusion criteria was adults aged 18 to 60 years of age with no ocular pathology. We excluded those with a history of previous ocular trauma or surgery. Those with a cup-disc ratio greater than 0.7, an intraocular pressure greater than 21 mm Hg, visual field abnormalities, a 1st-degree family member with glaucoma,

Table 1 Age, SE, axial length and 360° RNFL thickness in EM, LM, MM and HM groups

| Variables | EM | LM | MM | HM |
|-----------------------------|--------------|------------|-------------|------------|
| Age (y) | 31.85±10.55 | 31.10±9.64 | 31.38±11.03 | 31.15±8.68 |
| SE (D) | -0.11±0.21 | -1.47±0.73 | -4.45±0.81 | -8.25±1.41 |
| Axial length (mm) | 23.18±0.75 | 23.74±0.70 | 25.03±0.85 | 26.37±1.35 |
| Average RNFL thickness (µm) | 100.91±10.07 | 98.68±8.81 | 94.97±9.02 | 89.88±9.41 |

SD: Standard deviation; EM: Emmetropia; LM: Low myopia; MM: Moderate myopia; HM: High myopia.

Table 2 Peripapillary RNFL thickness and axial length in EM, LM, MM and HM groups

| Variable | EM (n=180) | LM (n=124) | MM (n=73) | HM (n=26) | ^a P | Post hoc Bonferroni correction |
|---------------------------------------|--------------|--------------|--------------|--------------|----------------|----------------------------------|
| 360° average RNFL thickness (µm) | 100.91±10.07 | 98.68±8.81 | 94.97±9.02 | 89.88±9.41 | <0.001 | EM>MM/HM LM>MM/HM |
| Superior quadrant RNFL thickness (µm) | 129.33±17.64 | 126.49±15.94 | 118.77±15.43 | 110.38±14.92 | <0.001 | EM>MM/HM LM>MM/HM |
| Inferior quadrant RNFL thickness (µm) | 134.82±17.64 | 129.31±16.91 | 121.56±15.90 | 109.92±15.15 | <0.001 | EM>LM/MM/HM LM>MM/HM MM>HM |
| Nasal quadrant RNFL thickness (µm) | 68.89±10.51 | 67.15±9.99 | 63.32±9.22 | 63.38±10.27 | <0.001 | EM>MM |
| Temporal quadrant RNFL thickness (µm) | 70.50±9.63 | 71.85±12.15 | 76.42±14.17 | 76.19±15.65 | <0.001 | EM<MM |
| Axial length (mm) | 23.18±0.75 | 23.74±0.84 | 25.03±0.85 | 26.37±1.35 | <0.001 | EM<LM/MM/HM LM<MM/HM MM<HM |

SD: Standard deviation; EM: Emmetropia; LM: Low myopia; MM: Moderate myopia; HM: High myopia. ^aANOVA was applied.

or peripapillary atrophy extending ≥ 17 mm from the center of optic disc were also excluded.

All subjects underwent a full ophthalmic examination, including distance Snellen visual acuity, refraction, slit-lamp biomicroscopy of the anterior segment, assessment of intraocular pressure by Goldmann applanation tonometry and dilated fundus examination. Myopic patients were classified based on their spherical equivalent (SE). The severity groups were as follows: low myopia (LM; SE greater than -0.5 D, up to -3.0 D), moderate myopia (MM; SE greater than -3.0 D, up to -6.0 D) and high myopia (HM; SE greater than -6.0 D)^[12]. The control group was emmetropia (EM), defined as a SE from +0.5 D to -0.5 D. Other examinations included visual field (SITA fast 24-2 Humphrey Field Analyser II, Carl Zeiss Meditec, Germany) and A-scan biometry (PAC SCAN 300 A Sonomed Digital Biomedic Ruler).

A Cirrus HD-OCT (Carl Zeiss Meditec, Inc. Germany) was used to measure the peripapillary RNFL thickness of both eyes of each subject. This was a spectral-domain OCT device with an acquisition rate of 27 000 A-scans per second. The optic disc cube 200×200 scan protocol was used to image the optic disc and the RNFL over the 6×6-mm² peripapillary region (200×200 data points). After the subject was properly seated, the iris was brought into view using the mouse-driven alignment system. The line scanning ophthalmoscopic image was focused, and the optic nerve head centered in the viewer. The software's automated built-in algorithms identified the center of the optic disc, and a circle measuring 3.46 mm in diameter was positioned automatically evenly around the disc center to

generate average, quadrant and clock-hour peripapillary RNFL measurements. A satisfactory scan required optic disc centration, images in clear focus and a signal strength of ≥ 6 . Images with movement artifact or signal strength of less than 7 were repeated once, if the second scan was also unusable, the eye was excluded from the study. The peripapillary RNFL parameters evaluated in this study consisted of mean 360°, superior, inferior, nasal and temporal quadrant thickness. Statistical analysis was performed using predictive analytics software (PASW) version 18.0. The mean peripapillary RNFL thickness between groups was compared using both analysis of variance (ANOVA) and analysis of covariance (ANCOVA) to control for potential confounders, with post hoc analysis using Bonferroni correction.

RESULTS

A total of 403 eyes of 403 subjects were included in this study. Approximately two-thirds were female (n=255). The mean age was 31.48±10.23y. There were 180 (44.7%) eyes with EM, 124 (30.8%) eyes with LM, 73 (18.1%) eyes with MM and 26 (6.5%) eyes with HM (Table 1).

There were significant inter-group differences for all RNFL parameters evaluated (Table 2). Myopic groups had a thinner average RNFL than the EM group, and their RNFL was observed to be thinner in all quadrants except temporally.

The myopic groups had a thinner average RNFL than the emmetropic group, with a statistically significant difference observed only between the HM and the EM group, after controlling for age, gender and axial length of the eye (P=0.017; Table 3).

Table 3 Comparison of average RNFL thickness between EM, LM, MM and HM groups

| Refractive error groups | Adjusted mean (95%CI) | ¹ Adjusted mean difference (95%CI) | ^a P |
|-------------------------|--|---|----------------|
| EM/LM | 99.22 (97.55, 100.90)/97.17 (95.30, 99.03) | 2.05 (-1.00, 5.11) | 0.450 |
| EM/MM | 99.22 (97.55, 100.90)/95.09 (92.45, 97.73) | 4.13 (-0.34, 8.61) | 0.088 |
| EM/HM | 99.22 (97.55, 100.90)/91.10 (86.41, 95.79) | 8.12 (0.96, 15.28) | 0.017 |
| LM/MM | 97.17 (95.30, 99.03)/95.09 (92.45, 97.73) | 2.08 (-2.06, 6.22) | 1.000 |
| LM/HM | 97.17 (95.30, 99.03)/91.10 (86.41, 95.79) | 6.07 (-0.60, 12.73) | 0.098 |
| MM/HM | 95.09 (92.45, 97.73)/91.10 (86.41, 95.79) | 3.99 (-2.02, 9.99) | 0.477 |

EM: Emmetropia; LM: Low myopia; MM: Moderate myopia; HM: High myopia; CI: Confidence interval. ¹ANCOVA was applied, with post-hoc Bonferroni correction; ^aP<0.05 is statistically significant.

Table 4 Comparison of mean inferior quadrant RNFL thickness between EM, LM, MM and HM groups

| Refractive error groups | Adjusted mean (95%CI) | ¹ Adjusted mean difference (95%CI) | ^a P |
|-------------------------|---|---|----------------|
| EM/LM | 130.73 (127.77, 133.70)/126.45 (123.15, 129.75) | 4.28 (-1.12, 9.69) | 0.217 |
| EM/MM | 130.73 (127.77, 133.70)/123.83 (119.16, 128.50) | 6.90 (-1.01, 14.82) | 0.127 |
| EM/HM | 130.73 (127.77, 133.70)/116.42 (108.13, 124.71) | 14.31 (1.65, 26.97) | 0.017 |
| LM/MM | 126.45 (123.15, 129.75)/123.83 (119.16, 128.50) | 2.62 (-4.70, 9.94) | 1.000 |
| LM/HM | 126.45 (123.15, 129.75)/116.42 (108.13, 124.71) | 10.03 (-1.76, 21.82) | 0.148 |
| MM/HM | 123.83 (119.16, 128.50)/116.42 (108.13, 124.71) | 7.41 (-3.22, 18.04) | 0.392 |

EM: Emmetropia; LM: Low myopia; MM: Moderate myopia; HM: High myopia; CI: Confidence interval. ¹ANCOVA was applied, with post-hoc Bonferroni correction; ^aP<0.05 is statistically significant.

The mean inferior quadrant RNFL was significantly thinner in the HM group compared to the EM group, after adjustment for age, gender and axial length ($P=0.017$; Table 4). There were no statistically significant differences of RNFL thickness among groups in the superior, nasal and temporal quadrants, after adjustment for confounders (data not shown).

DISCUSSION

The myopia epidemic has attracted growing concern worldwide, with researchers striving to identify risk factors for this condition^[13-16] as well as the optimal management of the disease and its complications^[17-19]. HM, especially, is associated with both maculopathy and glaucomatous optic neuropathy, rendering OCT an indispensable imaging modality in such patients^[20-21]. Our study provides mean RNFL values of Zeiss Cirrus HD-OCT as a baseline data for future assessment of myopic patients and demonstrates significant RNFL thickness differences between myopic patients and controls.

We observed that the mean RNFL thickness was significantly lower in highly myopic eyes compared to emmetropic eyes. This is in agreement with similar studies performed using RTVue-100 and Fourier domain OCT^[22-23]. Although increasing axial length and a more negative SE have both been associated with RNFL thinning^[24-25], we observed that the extent of this thinning was not statistically significant between the highly myopic group and the groups with low and MM. Our findings are in contrast to those of Sezgin Akcay *et al*^[26] and Kim *et al*^[27], who observed that patients with HM have a thinner average RNFL than those with low and moderate myopia.

Inter-group comparison of RNFL quadrant thickness revealed that the RNFL thinning seen in highly myopic eyes was not uniformly distributed; significant thinning was observed in the inferior quadrant RNFL of highly myopic eyes. However, our study failed to observe any significant inter-group differences in RNFL thickness of the superior and nasal quadrants. These findings are in inverse correlation to the two studies cited above^[26-27], which observed that the RNFL was thicker in the LM group than in the moderate and/or HM groups for the superior, nasal and inferior quadrants. The differences between our study and those cited may be due to the effect of confounders such as age^[28-29], as multivariate analysis was not applied in the aforementioned studies.

Retinal thinning in myopia has been attributed to reduced thickness of the middle to inner retina, which has been correlated functionally to reduced spatial resolution^[30]. This thinning has been explained by stretching of the ocular layers during eyeball elongation, as occurs in pathologic myopia^[31]. Increased axial length has also been associated with narrowed retinal arterioles^[32-33] and decreased peripapillary retinal flow perfusion, but whether this vascular compromise precedes or follows the RNFL thinning is still a matter of debate^[34].

Strengths of our study include a relatively large sample size, adjustment for confounders and the elimination of inter-observer errors by using a single operator to perform refraction, ocular biometry and OCT of the RNFL. However, we acknowledge our limitation of an unequal number of samples in each refractive error group, with the highly myopic group

comprising the smallest proportion. In addition, we could have inadvertently introduced selection bias when we excluded subjects with abnormal visual fields and increased cup-disc ratio. Finally, we omitted to adjust for magnification effect, which may also potentially affect the measured RNFL thickness^[35].

Despite our limitations, our study clearly demonstrates that highly myopic eyes have a thinner RNFL than normal eyes. On one hand, this thinning may be a risk factor for glaucoma development, as variations in the arrangement of optic nerve head fibers have been postulated to render myopic eyes more susceptible to glaucomatous damage^[5,36]. Of more clinical relevance, however, is that when evaluating peripapillary RNFL thinning in myopic eyes, the clinician must bear in mind that the age-matched normogram provided by the software to guide RNFL thickness assessment may not be valid, as it does not contain algorithms to adjust for axial length and refractive error.

Highly myopic eyes have thinner average RNFL and inferior quadrant RNFL compared to emmetropic eyes. Interpretation of RNFL thickness in highly myopic eyes should be performed with caution.

ACKNOWLEDGEMENTS

The authors would like to acknowledge Professor Dr. Syed Hatim Noor and Dr. Yee Siau Lin for their invaluable assistance in statistical analysis.

Conflicts of Interest: Tai ELM, None; Ling JL, None; Gan EH, None; Adil H, None; Wan-Hazabbah WH, None.

REFERENCES

- 1 Foster PJ, Jiang Y. Epidemiology of myopia. *Eye (Lond)* 2014;28(2):202-208.
- 2 Hopf S, Pfeiffer N. Epidemiology of myopia. *Ophthalmologe* 2017;114(1):20-23.
- 3 Wu PC, Huang HM, Yu HJ, Fang PC, Chen CT. Epidemiology of Myopia. *Asia Pac J Ophthalmol (Phila)* 2016;5(6):386-393.
- 4 Rose KA, French AN, Morgan IG. Environmental factors and myopia: paradoxes and prospects for prevention. *Asia Pac J Ophthalmol (Phila)* 2016;5(6):403-410.
- 5 Ma F, Dai J, Sun X. Progress in understanding the association between high myopia and primary open-angle glaucoma. *Clin Exp Ophthalmol* 2014;42(2):190-197.
- 6 Shim SH, Sung KR, Kim JM, Kim HT, Jeong J, Kim CY, Lee MY, Park KH; Korean Ophthalmological Society. The prevalence of open-angle glaucoma by age in myopia: the Korea National Health and Nutrition Examination Survey. *Curr Eye Res* 2016;42(1):1-7.
- 7 Hsu CH, Chen RI, Lin SC. Myopia and glaucoma: sorting out the difference. *Curr Opin Ophthalmol* 2015;26(2):90-95.
- 8 Fu L, Aspinall P, Bennett G, Magidson J, Tatham AJ. The influence of optical coherence tomography measurements of retinal nerve fiber layer on decision-making in glaucoma diagnosis. *Curr Eye Res* 2017;42(4):575-582.
- 9 Hua Z, Fang Q, Sha X, Yang R, Hong Z. Role of retinal nerve fiber layer thickness and optic disk measurement by OCT on early diagnosis of glaucoma. *Eye Sci* 2015;30(1):7-12.
- 10 Kotowski J, Wollstein G, Ishikawa H, Schuman JS. Imaging of the optic nerve and retinal nerve fiber layer: an essential part of glaucoma diagnosis and monitoring. *Surv Ophthalmol* 2014;59(4):458-467.
- 11 Zangwill LM, Bowd C. Retinal nerve fiber layer analysis in the diagnosis of glaucoma. *Curr Opin Ophthalmol* 2006;17(2):120-131.
- 12 Cline D, Hofstetter HW, Griffin JR. Dictionary of Visual Science. 4th ed: Butterworth-Heinemann; 1997.
- 13 Cuellar-Partida G, Lu Y, Kho PF, Hewitt AW, Wichmann HE, Yazar S, Stambolian D, Bailey-Wilson JE, Wojciechowski R, Wang JJ, Mitchell P, Mackey DA, MacGregor S. Assessing the genetic predisposition of education on myopia: a mendelian randomization study. *Genet Epidemiol* 2016;40(1):66-72.
- 14 Gong B, Qu C, Huang XF, Ye ZM, Zhang DD, Shi Y, Chen R, Liu YP, Shuai P. Genetic association of COL1A1 polymorphisms with high myopia in Asian population: a Meta-analysis. *Int J Ophthalmol* 2016;9(8):1187-1193.
- 15 Guggenheim JA, Williams C; UK Biobank Eye and Vision Consortium. Childhood febrile illness and the risk of myopia in UK Biobank participants. *Eye (Lond)* 2016;30(4):608-614.
- 16 Parssinen O, Kauppinen M. Associations of reading posture, gaze angle and reading distance with myopia and myopic progression. *Acta Ophthalmol* 2016;94(8):775-779.
- 17 Davidson SL, O'Hara M, Wagner RS. Management of progressive myopia. *J Pediatr Ophthalmol Strabismus* 2016;53(3):134-136.
- 18 Huang J, Wen D, Wang Q, et al. Efficacy comparison of 16 interventions for myopia control in children: a network Meta-analysis. *Ophthalmology* 2016;123(4):697-708.
- 19 Wolffsohn JS, Calossi A, Cho P, et al. Global trends in myopia management attitudes and strategies in clinical practice. *Cont Lens Anterior Eye* 2016;39(2):106-116.
- 20 Chen LW, Lan YW, Hsieh JW. The optic nerve head in primary open-angle glaucoma eyes with high myopia: characteristics and association with visual field defects. *J Glaucoma* 2016;25(6):e569-e575.
- 21 Ohno-Matsui K. Pathologic myopia. *Asia Pac J Ophthalmol (Phila)* 2016;5(6):415-423.
- 22 Kita Y, Kita R, Takeyama A, Tomita G, Goldberg I. Effect of high myopia on glaucoma diagnostic parameters measured with optical coherence tomography. *Clin Exp Ophthalmol* 2014;42(8):722-728.
- 23 Malakar M, Askari SN, Ashraf H, Waris A, Ahuja A, Asghar A. Optical coherence tomography assisted retinal nerve fibre layer thickness profile in high myopia. *J Clin Diagn Res* 2015;9(2):NC01-3.
- 24 Leung CK, Mohamed S, Leung KS, Cheung CY, Chan SL, Cheng DK, Lee AK, Leung GY, Rao SK, Lam DS. Retinal nerve fiber layer measurements in myopia: an optical coherence tomography study. *Invest Ophthalmol Vis Sci* 2006;47(12):5171-5176.
- 25 Zhao Z, Jiang C. Effect of myopia on ganglion cell complex and peripapillary retinal nerve fibre layer measurements: a Fourier-domain

RNFL thickness in myopia and controls

- optical coherence tomography study of young Chinese persons. *Clin Exp Ophthalmol* 2013;41(6):561-566.
- 26 Sezgin Akcay BI, Gunay BO, Kardes E, Unlu C, Ergin A. Evaluation of the ganglion cell complex and retinal nerve fiber layer in low, moderate, and high myopia: a study by RTVue spectral domain optical coherence tomography. *Semin Ophthalmol* 2017;32(6):682-688.
- 27 Kim MJ, Lee EJ, Kim TW. Peripapillary retinal nerve fibre layer thickness profile in subjects with myopia measured using the Stratus optical coherence tomography. *Br J Ophthalmol* 2010;94(1):115-120.
- 28 Uchida H, Yamamoto T, Araie M, Tomita G, Shirakashi M, Yoshikawa K; HRT Study Group. Topographic characteristics of the optic nerve head measured with scanning laser tomography in normal Japanese subjects. *Jpn J Ophthalmol* 2005;49(6):469-476.
- 29 Ozdek SC, Onol M, Gürelik G, Hasanreisoglu B. Scanning laser polarimetry in normal subjects and patients with myopia. *Br J Ophthalmol* 2000;84(3):264-267.
- 30 Wolsley CJ, Saunders KJ, Silvestri G, Anderson RS. Investigation of changes in the myopic retina using multifocal electroretinograms, optical coherence tomography and peripheral resolution acuity. *Vision Res* 2008;48(14):1554-1561.
- 31 Yanoff M, Fine BS. *Ocular Pathology: A Text and Atlas*. Philadelphia: Harper & Row 1982.
- 32 Gopinath B, Wang JJ, Kifley A, Tan AG, Wong TY, Mitchell P. The association between ocular biometry and retinal vascular caliber is comparable from early childhood to adolescence. *Invest Ophthalmol Vis Sci* 2013;54(2):1501-1508.
- 33 Tai EL, Li LJ, Wan-Hazabbah WH, Wong TY, Shatriah I. Effect of axial eye length on retinal vessel parameters in 6 to 12-year-old Malay girls. *PLoS One* 2017;12(1):e0170014.
- 34 Wang X, Kong X, Jiang C, Li M, Yu J, Sun X. Is the peripapillary retinal perfusion related to myopia in healthy eyes? A prospective comparative study. *BMJ Open* 2016;6(3):e010791.
- 35 Bae SH, Kang SH, Feng CS, Park J, Jeong JH, Yi K. Influence of myopia on size of optic nerve head and retinal nerve fiber layer thickness measured by spectral domain optical coherence tomography. *Korean J Ophthalmol* 2016;30(5):335-343.
- 36 Cahane M, Bartov E. Axial length and scleral thickness effect on susceptibility to glaucomatous damage: a theoretical model implementing Leplace's law. *Ophthalmic Res* 1992;24(5):280-284.